BRL



CONTRACT 169 REPORT NO. 5

UPPER ATMOSPHERE WINDS FROM

GUN LAUNCHED VERTICAL PROBES

(Barbados, 17-25 February 1966)

SPACE INSTRUMENTS RESEARCH, INC.

DISCLAIMER NOTICE

THIS DOCUMENT IS BEST QUALITY PRACTICABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

UPPER ATMOSPHERE WINDS FROM
GUN LAUNCHED VERTICAL PROBES
(Barbados February 17-25 1966)

Prepared for

U. S. Army Ballistic Research Laborato ies Aberdeen Proving Ground, Maryland

Contract No. DA-01-009-AMC-169(X)

Tas docueros is unimited. its distribution is unimited.

Prepared by:

Sobert & Teller

Robert N. Fuller Research Physicist Approved by:

Howard D. Edwards

Technical Director

Space Instruments Research, Inc. Atlanta, Georgia

January 1967

Dr

DEC 18 1957

س سا الماس

TABLE OF CONTENTS

	Page
Introduction	1
Data Acquisition	2
Data Reduction	4
Interpretation of Data	6
Illustrations	8
Synopsis of Results	10
References	11
Table of Trail Information	14
WIND PROFILES:	
Eight Trail Releases February 17-25, 1966	15

INTRODUCTION

A continuing study of upper atmospheric winds over the lower
West Indies has been made possible by the firing of high altitude
bailistic probes from a sixteen-inch gun located on the Island of
Barbados. These firings are being carried out by the U. S. Army
Ballistics Research Laboratories, Aberdeen Proving Ground, Maryland,
under the direction of Dr. Charles H. Murphy, and by the Space Research
Institute of McGill University, Canada, under the direction of Dr. G.V.
Bull.

Atmospheric winds are studied by releasing chemical trails from the gun-fired probes during the upper portion of their trajectories. To date, the primary chemical which has been released is trimethyl aluminum (TMA). TMA produces a chemiluminescent glow in regions of the atmosphere above 85 kilometers, thus allowing the trails to be photographed while being distorted by upper atmosphere winds. The photographs are then reduced to provide wind information by Space instruments Research, inc. (SIR), using computer techniques.

The purpose of this report is to summarize results of these studies for the period from February 15 through February 25, 1966. Previous results for winds over Barbados, West Indies, and Yuma, Arizona, are covered in Technical Reports Nos. 1, 2, 3, and 4.

DATA ACQUISITION

The chemical trails are formed almost vertically over the Island of Barbados (longitude 59.4°W, latitude 13.0°N) and extend from an altitude of 85 kilometers through apogee. In some firings, TMA is also released on the down leg of the trajectory. To the unaided eye, the chemical release first appears as a straight white trail resembling a jet contrail. Within a minute or so, the trail is distorted into strainge shapes by the upper atmospheric winds (see Fig. 1) and fades from view within approximately fifteen minutes after initial release.

Space instruments Research has established eight photographic triangulation stations on the Islands of Barbados, St. Vincent, Grenada, and Tobago, with two sites per Island. These latter islands are located to the west and south of Barbados at distances of 190 to 290 kilometers (see Fig. 2). While only one site on each of two Islands is required for data reduction purposes, the eight sites have been found necessary because of cloud conditions in the area.

Equipment at each site, built by SIR, consists of a camera unit containing two seven-inch focal length cameras mounted on a concrete pedestal, and an electronic control. Cameras are automatically pulsed to take exposures of 3, 6, and 12 seconds duration every 30 seconds.

Since commercial power is either unreliable or unavailable at many site locations, SIR has developed a battery operated i15-volt power supply for the control equipment. The power supply is tuning-fork controlled and provides 60 cycle power with an accuracy of 0.005%

for the camera programmer so that pictures can be taken simultaneously at each site. A data block containing 24 tiny lights, mounted in each camera unit, records time, firing number, and site information in the corner of each frame of film.

A short wave radio net connecting all sites and the launch control center has been installed by SIR to enable the launch control officer on Barbados to be informed of weather conditions on the islands and to synchronize picture-taking operations with the firing of the gun. Most sites are operated by local personnel who have been trained by SIR.

During a typical night's operation, the gun is fired at one to two-hour intervals, from sunset to sunrise. Photographs are taken by all sites during the time that the trail is visible. The film is returned to Atlanta for processing and data reduction.

DATA REDUCTION

Several computer programs have been developed which make it possible to calculate upper atmosphere winds from measurements made directly on the photographs of the luminous trails.

Since the method used is basically three-dimensional triangulation using spherical trigonometry, it is necessary to know precisely the direction each camera was pointed during a given firing. The direction is determined by first taking accurate measurements of the locations of several star images on the film, and then computing the azimuth and elevation of the optical axis of the camera by means of a computer program. This computer program makes use of the celestial coordinates of some 6,000 stars which have been stored on magnetic tape.

Wind speeds and directions are then determined from the location of the trail in space at a succession of known times. The location is found, using either a point location program or a trail location program, or both, and depends on the physical shape of the chemical release cloud.

Point location method. If the chemical release exhibits discrete points (resulting either from turbulence or from the nature of the release mechanism) and these points can be identified on films from two or more sites, the point location program can be used to calculate the position of each point in longitude, latitude, and altitude above sea level.

These calculations mare made from data taken at successive times.

A wind program is then used to calculate both vertical and horizontal and from the motion of these points as a function of time.

Trail location method. Most of the chemical releases produce a smooth trail having few, if any, identifiable points. In such cases, film coordinates of a large number of incremental points along the film image of the trail are fed into the computer from data from two or more sites. The trail location program attempts to triangulate each point from one site with many points from another site, finally choosing points from both sites whose optical paths from camera into space form the closest spatial intersection. After doing many hundreds of such calculations, the computer is able to construct coordinates for a mathematical curve in the shape of the trail in space. Then, as with the point location program, winds can be determined from the motion of the curva with time. Here, however, it must be assumed that vertical winds are essentially zero. This assumption is borne out by previous studies which have shown vertical winds in this altitude region to be of the order of a few meters per second compared to horizontal winds ranging up to 150 meters per second.

Corrections for variables such as atmospheric refraction, rotation of camera about optical axis, and camera focal length, are incorporated into the programs to maintain high accuracy. Focal length and camera rotation are, in fact, calculated from measurements of the positions of star images on the films.

INTERPRETATION OF DATA

In the remainder of this report, horizontal wind velocities are presented in tabular form and in plots of wind speed, direction, and components.

Winds were calculated at altitude intervals of one kilometer. Points on the various plots show the actual computed result, as listed in the table preceding the plot. A curve has been fitted to each set of points to aid in detecting wind patterns and to indicate reliability of the plotted results. Each curve has been drawn with a knowledge of intermediate results leading to the wind calculations and of the consistency of the winds as calculated between each of the five or more time intervals used. In cases where point-to-point curve fitting was not thought to reflect actual variations in wind speed, direction, or components, a more appropriate smooth curve has been drawn Otherwise, the curves are fitted directly to the data points. Results of certain portions of the trails are at times less accurate than others due to the spatial orientation of those trail segments relative to the available photographic stations. Less accurate data also can result from photographs obscured by haze and clouds and from trails of short duration.

Wind speed plot. This plot shows the speed of the wind in meters per second as a function of height in kilometers above sea level.

Wind direction plot. The wind vector is considered to point in the direction toward which the wind is moving. The direction plot shows the direction of this vector in degrees clockwise from north

as seen from above. Thus, a wind direction toward the east would be 90 degrees.


Wind components plot. While plots of wind direction and speed do completely describe the wind vector, it has been found helpful in studying wind patterns to present the north-south and east-west velocity components of the vector. In the north-south plot, north is positive; south is negative. In the east-west plot, east is positive, west negative. Components are plotted in meters per second versus height in kilometers.

The wind direction and components described above are referenced to true north. In addition, components have been calculated relative to magnetic north for comparison with other ionospheric phenomena. These components are not plotted but are listed in the tabulations preceding each set of plots.


Fig. 1

Photographs of firing St. Kitts

Photographs taken 167 seconds after firing:








Barbados


St. Vincent

Tobago

These pictures were taken from three islands just as the vehicle reached apogee. Note that the winds have already distorted the trail. Numbers indicate altitude in Kilometers.

Photographs taken 250 seconds after firing:






Barbados

Tobago

This set of pictures was taken at completion of the downtrail. The uptrail shows the continued effect of the winds, while the downtrail is new and only slightly distorted. Stars can be seen in the background of these pictures. The positions of these stars are used to determine the exact direction each camera was aimed.

Fig. 2

Location of S.I.R. photographic stations
H.A.R.P.-Barbados



Two stations are located on each of the four islands, as shown. While only one station on each of any two islands is sufficient for determination of winds by triangulation, several stations were found necessary because of prevalent cloud conditions in the area. Accuracy of the data reduction is also increased by use of films from more than two Islands.

SYNOPSIS OF RESULTS

The following comments may be helpful in interpreting the data contained in this report. Only those shots with unusual characteristics are discussed.

INAUGUA

Downtrail results above 111 kilometers were significantly different from uptrail results. Two curves are shown in this area. Below this region it was not clear whether or not there were actual differences between the up and down results, so only one curve was drawn.

ST. KITTS

The region 95 through 98 kilometers gave poor results. The points are not thought to be accurate, thus a smooth curve was drawn through them.

MONTSERRAT

Very poor photographic data due to bad weather made analysis of this trail difficult. Only a small portion of the trail length was useable. Winds shown are generally less accurate than other trails.

ST. THOMAS

This trail had a very unusual characteristic. Above 110 kilometers, up through apogee, and into the downtrail, the release split into two crails. These two trails showed different apogees and different winds.

Photographs of this release seem to indicate that the two trails rejoined in the down leg, but for two reasons we believe this is not the case. First, triangulation results show different positions for the two trails near where they appear to join. Second, three other that school data has not set been reduced show very similar double trails, and on these the downtrails do not rejoin. In fact, the lower trails seem to join the "reentry bags" -- indicating that the second that may be formed by the vaporization of previously frazen TMA or stalls as the, follow their own trajectory. We believe the lower trail of this boot twisted in front of the upper, causing them to appear to join.


Deta from the up leg of this second trail was too poor to give useful results. Thus results for one uptrail and two downtrails are shown. The lower (reviporization) trail is called trail "A".

Figure three shows a ground projection of these trails.

Fig. 3

GROUND PLOT

TRAIL NO. 49 ST. THOMAS T+263 Seconds



REFERENCES

- 1. Albritton, C. L., L. C. Young, H. D. Edwards, and J. L. Brown, "Fosition Determination of Artificial Clouds in the Upper Atmosphere," Photogrammetric Engineering, September 1962.
- 2. Armstrong, f. 8, "Observations of Luminous Clouds Produced in the Upper Atmosphere by Exploring Grenades 1, 11 and 111," Planet, Space Sci., 11, 737-758, 1363.
- 3. Bedinger, J. F., Compendium of Wind Data from the Vapor Trail Technique," GCA Tech. Report. 66-7-N, March 1966.
- 4. Blamont, J. E., "Turbulence in Atmospheric Motions Between 90 and 130 km of Altitude," Planetary and Space Sciences, 10, 89-101, 1963.
- 5. Bull, G. V., C. H. Murphy, "Gun Launched Missiles for Upper Atmosphere Research," AIAA Preprint No. 64-18, January 1964.
- Bull, G. V., H. J. Luckert, "Report of the March 1965 Test Firing Series." Project HARP, McGill University Report SRI-H-R-9, July 1965.
- 7. Cato, O., "Turbulent Diffusion of Sodium Vapor Trails in the Upper Atmosphere," GCA Technical Report No. 65-5-N, Contract NASw-1083, March 1965.
- 8. Champion, K. S. W., "Atmospheric Structure and Its Variations in the Lower Thermosphere," <u>Planet</u>, <u>Space Sci.</u>, 13, 325-338, 1965.
- Edwards, H. D., M. M. Cooksey, C. G. Justus, R. N. Fuller, D. L. Albritton, N. W. Rosenberg, "Upper Atmosphere Wind Measurements Determined from Twelve Rocket Experiments," J. Geophys. Res., 68, 3021-3032, 1963.
- Edwards, H. D., C. G. Justus, D. C. Kurts, "Evening Twilight Winds from 68 to 140 Kilometers for May 21, 1963," J. Geophys. Res., 68, 6062-6063, 1963.
- il. Elford, W. G., and R. G. Roper, "Turbulence in the Lower Thermosphere," to be published in <u>Space Research</u> VII, 1967.
- 12. Golomb, D., and M. M. MacLeod, "Diffusion Coefficients in the Upper Atmosphere fro Chemiluminous Trails," J. Geophys. Res., 71, 2299-2305, 1966.
- Golomb, D., N. W. Rosenberg, C. Abaronian, J. A. Hill, and H. L. Alden, "Oxygen Atom Determination in the Upper Atmosphere by Chemiluminescence of vitric Oxide," <u>J. Geophys. Res.</u>, 70, 1155-1173, 1965.
- 14. Hines, C. O., "Ionospheric Movements and Irregularities," Research in Geophysics, Vol. 1, 299-318, 1964.

- 15. Hines, C. O., 'Minimum Vertical Scale Sizes in the Wind Structure Above 100 Kilometers," J. Geophys. Res., 69, 2847-2848, 1964.
- 16. Hines, C.O., 'Dynamical heating in the Upper Atmosphere," J. Geophys. Res., 70, 177-183, 1965
- 17. Hines, C. O., "Di. rn. 1 Time in the Upper Atmosphere," Jour. Geophys. Research 71, 1453, 1466.
- 18. Hines, C. O., 'Or the Analysis and Interpretation of Winds Observed at Heights of 85 to 135 Kilometers: A Rebuttal," J. Geophys. Res. 71, 1461, 1966.
- Johnson, E. R., and R. H. Lloyd, "Determination of Diffus in Coefficients from Observations on Grenade Glow Clouds," <u>Australian Jour. Phys.</u> 16, 490-499, 1963.
- 20. Justus, C. G., "The Energy Balance of Turbulence in the Upper Atmosphere," J. Geophys. Res. 71, No. 15, August 1, 1966.
- 21. Justus, C. G., H. D. Edwards, R. N. Fuller, "A Method Employing Star Backgrounds for Improving the Accuracy of the Location of Clouds or Objects in Space," Photogrammetric Engineering, July 1964.
- 22. Kochanski, A., "Atmospheric Motions from Sodium Cloud Drifts at Four Locations," Monthly Weather Review, Vol. 94, No. 4, April 1966.
- 23. Lloyd, K. H., and L. M. Shappard, "Atmospheric Structure at 130-200 km Altitude from Observations on Grenade Glow Clouds During 1962-63," Australian Jour. Phys., 19, 323-342, 1966.
- 24. Morgan, A. W., 'Measurements of Winds by Chemical Releases in the Upper Atmosphere, NASA Technical Memorandum, NASA TM X-53363, December 1965.
- 25. Murphy, C. H., G. V. Bull, H. D. Edwards, "Upper Atmosphere Winds Measured by Gun Launched Projectiles," AMS/AIAA Conference on Aerospace Mcteorology, March 1966, and J. Geophys. Res. (in press).
- 26. Noel, T. M., "A Measurement of Turbulence Power and Small Eddy Scale Near 105 Kilometers," J. Geophys. Res., 68, 2862-2863, 1963.
- 27. Nordberg, W., "Rocket Soundings in the Mesosphere," in NASA SP-49 (Meteorological Observations above 30 km) 1964.
- 28. Roper, R. G., "Dissipation of Wind Energy in the Height Range 80 to 140 km," Jour. Geophys. Res., 71, September 15, 1966.
- 29. Roper, R. G., "The Semidiurnal Tide in the Lower Thermosphere." Jour. Geophys. Res. (accepted for publication probably Dec. 15, 1966).

- 30. Roper, R. G., 'Atmospheric Turbulence in the Meteor Region," <u>Jour.</u> Geophys. Res. (accepted for publication, probably December 15, 1966.)
- Fosenberg, N. W., H. D. Edwards, and J. W. Wright, "Ionospheric Winds: Motions into Night and Sporadic E Correlations," <u>Space Research</u> 4, 171-181, 1964.
- 32. Rosenberg, N. W., D. Golomb, E. F. Allen, "Chemiluminescence of Trimethyl aluminum Released into the Upper Atmosphere," J. Geophys. Res. 68, 5895-5898, 1963.
- 33. Resemberg, N. W., D. Golomb, E. F. Allen, "Resonance Radiation of Al0 from Trimethyl Aluminum Released into the Upper Atmosphere," J. Geophys. Res. 69, 1451-1454, 1964.
- 34. Rosenberg, N. W., D. Golomb, E. F. Allen, "Chemiluminescent Techniques for Studying Nighttime Winds in the Upper Atmosphere," J. Geophys. Res. 68, 3328-3330, 1963.
- 35. Rosenberg, N. W., C. G. Justus, "Space and Time Correlations of Ionospheric Winds," Radio Science, 1, No. 2, February 1966.
- 36. Shappard, L. M., and K. H. Lloyd, "Atmospheric Density and the Diffusion of Grenade Glow Clouds," Planet Space Sci., 12, 317-318, 1964.
- 37. Zimmerman, "Small-Scale Wind Structure Above 100 km," Jour. Geophys. Res., 69, 784-785, 1964.

TABLE OF TRAIL INFORMATION

TRAIL NO.	NAME	DATE	TIME (A.S.T.)	ALTITUDES (KM.)
в43	Inaugua	17 February, 1966	21:03:00	92 - 123
B44	St. Kitts	23 Feb.uary, 1966	20:46:00	87 - 131
845	St. Lucia	23 February, 1966	22:03:00	95 - 117
B46	Montserrat	23 Pebruary, 1966	23:21:00	111 - 120
B47	Nevis	24 Pebruary, 1966	00:25:00	91 - 122
B48	Puerto Rico	24 February, 1966	03:27:00	90 - 123
B49	St. Thomas	24 February, 1966	05:23:00	96 - 118
в50	Plamingo	25 February, 1966	18:43:00	94 - 130

TABULATIONS AND PLOTS


Eight Trail Releases - February 17-25, 1966


	WIND	WIND	WIND COMPONENTS (M/S)			
ALTITUĐE	HEADING	VELOCITY		RAPHIC		
(KM)	(DEG)	(M/S)	N-S	E-W	N-S	E-W
92.0	299.8	97.9	48.7	-85.0	64.9	-73.4
93.0	312.0	112.3	75.2	-83.5	90.5	-66.6
94.0	324.7	111.4	90.9	-64.4	102.0	-44.7
95.0	337.4	112.7	104.0	-43.4	110.6	-21.5
96.0	345.6	117.0	113.3	-29.0	116.8	-5.5
97.0	345.5	136.8	132.4	-34.1	136.6	-6.7
98.0	344.2	143.9	138.5	-39.1	143.5	-10.3
99.0	344.4	143.5	138.2	-38.7	143.2	-10.0
100.0	353.7	113.7	113.0	-12.5	113.2	10.6
101.0	6 • 4	105.7	105.0	11.8	100.5	32.8
192.0	13.7	100.2	97.3	23.8	90.5	43.0
103.0	23.0	75.2	69.2	29.4	61.8	42.8
194.0	31.0	62.8	53.8	32.4	46.1	42.6
105.0	44 • 2	44.2	31.7	30.8	24 • 8	36.6
166.0	105.0	21.6	-5.6	20.9	-9.7	19.3
197.0	161.4	17.3	-16.4	5•5	-17.2	2.1
108.0	248 . 5	22.6	-8.3	-21.0	-3.9	-22.2
109.0	265.1	14.9	-1.3	-14.8	1.7	-14.8
110.0	336.6	18.1	16.6	-7.2	17.7	-3.7
111.0	43.5	20.2	14.6	13.9	11.5	16.6
112.0	49.0	35.9	23.5	27.1	17.5	31.3
113.0	63.6	53.6	23.8	48.0	13.6	51.8
114.9	68.9	66•4	23.9	62.0	10.9	65.5
115.0	69.8	72.2	24.9	67.8	10.7	71.4
116.0	71.2	79.5	25.7	75.3	10.0	78.9
117.0	71.9	87.9	27.3	83.5	9•9	87.3
118.0	77.6	87.0	18.7	85.0	1 • 2	87.0
119.0	88.1	81.8	2.7	81.7	-13.9	80.6
120.0	96•4	76.6	-8.5	76.1	-23.7	72.8
121.0	99.8	74.9	-12.7	73.8	-27.3	69.7
122.0	162.7	75.1	-16.5	73.3	-31.0	68.5
123.0	101.4	73.8	-14.6	72.3	-28.9	67.9


TRAIL NO. B43 INAUGUA 17 FEBRUARY 1966 21-03-00 AST

	WIND	WIND	1	WIND COMPO	NENTS (M/S)
ALTITUDE	HEADING	VELOCITY	GEOGRAPHIC		MAGNETIC	
(KM)	(DEG)	(M/S)	N-S	E-w	N-5	E-4
107.0	212.4	7.5	-6.3	-4.0	-5.4	-5.2
108.0	221.6	20.9	-15.6	-13.9	-12.5	-16.8
169.0	261.7	19.7	-2 • 8	-19.5	1.2	-19.7
110.0	328.7	16.2	13.8	-8.4	15.2	- <u>r</u> . 4
111.0	19.0	16.7	15.8	5.5	14.4	8.6
112.0	62.4	39.6	18.3	35.1	10.8	38.1
113.0	72.2	59.4	18.2	56.6	6.4	59.1
114.0	76.2	67.0	16.0	65.1	2.5	67.0
115.0	77.3	86.9	19.2	84.8	1.7	36.9
116.0	78•4	99.8	20.0	97.7	-0.1	99.7
117.0	80.9	101.6	16.0	100.3	-4.6	101.5
118.0	64•2	100.6	10.1	100.1	-10.3	100.1
119.0	91.4	93.7	-2.2	93.7	-21 • 1	91.3
120.0	94 • 8	94.4	-8.0	94.1	-26.8	90.5
121.0	97.3	97.5	-12.3	96.7	-31.6	92.2


BARBADOS DOWN TRAIL










	WIND	WIND		WIND COURS	NCNTS ////	
ALTITUDE	HEADING	VELOCITY	GEOC	WIND COMPO RAPHIC		
(KM)	(DEG)	(M/S)	N-S			NETIC
87.0	195.8	92.5	-89.0	E-W	N-S	E- *
88.0	197.7	66.9	-63.7	-25 • 1	-82 • 1	-42.6
89.0	206.5	79.7	-71.4	-20 • 4	-58.3	-32.8
90.0	210.2	74.7	-64.6	-35 • 5	-62.8	-49.7
91.0	229•2	61.4	-40.2	-37.6	-55.7	-49.9
92.0	224.1	60.3	-43.4	-46 • 5	-30.0	-53.7
93.0	234.6	62.2	-36.1	-42.0 -50.7	-34.0	-49.9
94.0	265.7	81.5	-6.1		-25.1	-56.9
95.0	297.5	78.9	36.5	-81.2 -70.0	10.4	-80.8
96.0	296 • 1	69 . 8	30.7	-70 . 0	49.9	-61.2
97.0	304.5	62.1	35.2	-62 • 7	42.7	-55.2
98.0	311.5	85.3		-51.1	44.8	-42.9
99.0	316.7	85.0	56.5	-63 . 9	68.2	-51.2
100.0	321.9	82.6	61.8	-58.3	72.3	-44.6
101.0	327.1	79.7	65.0	-50.9	73.9	-36.7
102.0	328.2	87.6	66.9	-43.2	74.2	-28.8
193.0	323.5	103.4	74.4	-46 • 2	82.2	-30.2
104.0	323.1	109.8	83.2	-61.5	93.9	-43.4
195.0	323.3	114.1	87.7	-65.9	99.2	-46.8
106.0	316.4	95.6	91.5	-68.2	103.4	-48.3
197.9	263.2		69.2	-66.0	81.1	-50.7
108.0	262.1	41.0	-4.9	-40.7	3.4	-40.9
199.0	278•7	28,8 15.0	-4.0	-28.5	1.8	-28.7
110.0	18.0	10.4	2 • 3	-14.8	5.2	-14.0
111.0	55.1		9•8	3 • 2	9.0	5.1
112.0	57.5	28.6 39.8	16.3	23.5	11.2	26.3
113.0	64 • 1	42.7	21.4	33.6	14.2	37.2
114.0	73.5	41.7	18.7	38 • 4	10.6	41.4
115.0	79.3	82.5	11.9	40.0	3.6	41.6
116.0	78.5	88.7	15.2	81.1	-1.5	82.5
117.0	77.3	90.2	17.6	87.0	-0.3	88.8
118.0	76.4	91.5	19.9	88.0	1.7	90.2
119.0	76.9	89.3	21.5 20.3	88.9	3.1	91.4
120.0	82.2	80.8		87.0	2.3	89.1
121.0			11.0	80.0	-5.4	80.6
122.0	83•7 84•4	76•4	8 • 3	76.0	-7.2	76.1
123.0	83.5	76.6 77.1	7•5 8•7	76 • 3	-8 • 1	76.2
124.0	82.5	77.6	10.1	76•7	-7.0 5.7	76.9
125.0	85.8	70.9	5.1	77•0	-5.7	77.5
126.0	91.1	62.1	-1.2	70•7 62•0	-9.3	70.3
127.0	92.9	57.7	-3.0		-13.7	60.5
128.9	95.3	53.0	-3.0 -4.9	57•6	-14.6	55.8
129.9	96•1	49.5	-4·9 -5·3	52 • 8	-15.5	50.7
130.9	104.7	43.4	-11.0	49.3	-15.1	47.2
131.0	119.9	35.1	17.5	42 • 0	-19.3	38.9
	, • ,	J J = 1	1100	30•4	-23.3	26.:






	WIND	WIND	WIND COMPONENTS (M/S)				
ALTITUDE	HEADING	VELOCITY		RAPHIC	MAGNETIC		
(KM)	(DEG)	(M/S)	N-S	E-W	N-S	E-w	
95.0	300.6	78.4	39.9	-67.5	52.7	-58.1	
96.0	308.6	86.4	53.9	-67.5	66 • 4	-55.2	
97.0	318.6	90.6	68.0	-59.9	78.7	-44.9	
98.0	322 • 1	92.9	73.2	-57.1	83.2	-41.1	
99.0	321.9	95.5	75.2	-58.9	85.5	-42.5	
100.4	321.7	95.0	74.6	~ 58∙\$	85.0	-42.6	
101.0	323.6	93.9	75.6	-55•7	85.3	-39.3	
102.0	319.5	90.6	68.9	~58.9	79.4	-43.8	
103.0	313.0	89.4	60.9	-65.4	72.9	-51.8	
104.0	290•8	55.4	19.7	-51.8	29.8	-46.8	
105.0	244•6	39.7	-17.0	-35.9	-9.4	-38.6	
106.0	229.5	34.6	-22.5	-26.3	-16.7	-30.3	
197.0	209•1	30.3	-26.4	-14.7	-22.9	-19.7	
198.0	168.4	26.3	-25.8	5•3	-26.3	0.3	
109.0	140.3	38.9	-29.9	24.9	-34.3	19.3	
110.0	126.1	53.8	-31.7	43.5	-39.8	36.2	
111.0	117.4	68.8	-31.7	61.1	-43.4	53.4	
112.0	120.4	69.2	-35.0	59.7	-46 • 3	51.4	
113.0	115.4	69.1	-29.6	62.4	-41.6	55.1	
114.0	117.1	65.4	-29.8	58•2	-40.9	51.0	
115.0	115.3	62.2	-26.6	56.2	-37.4	49.7	
116.0	99•9	62.7	-10.8	61.8	-23.1	58.3	
117.0	87.0	77.3	4 • 0	77•2	-11.7	76.4	










TRAIL NO. B46 MONTSERRAT
23 FEBRUARY 1966 23-21-00 AST

BARBADOS


	WIND	WIND	h	IND COMPO	NENTS (M/S)	
ALTITUDE	HEADING	VELOCITY	GEOGR	APHIC	MAGN	ETIC
(KM)	(DEG)	(M/S)	N-S	E-W	N-5	E-₩
111.0	128•9	57.4	-36.0	44•7	-44.3	36•5
112.0	124•7	84.8	-48 • 3	69•7	-61 • 4	58.5
113.0	125•2	91.2	-52 • 6	74•5	-66 • 6	62.3
.14.0	80•2	51.0	8 • 7	50.3	-1.6	5i.u
1.5.0	59•6	48.6	24 • 5	41.9	15.6	46.û
116.0	45 • 2	54.8	38•6	38.9	29•9	45.9
	67 • 9	46.7	17•6	43.3	8•5	45.0
118.0	105 • 8	30.2	-8.2	29 • 1	-13.9	26.8
	91 • 2	31.5	-0.7	31 • 4	-7.0	30.6
120.0	50.3	23.4	15.0	18.0	11.1	20.7









	WIND	WIND			NENTS (M/S)
ALTITUDE	HEADING	VELOCITY	GEOGRAPHIC		MAGI	ETIC
(KM)	(DEG)	(M/S)	N-S	E-W	N-S	[- W
90	250.5	64.4	-21.5	-60.7	-8.8	-63.8
90	261.0	80.0	-12.6	-79.0	3.6	-74.9
93 .0	279.1	76.1	12.1	-75 · i	27.0	-71.1
94.0	294.2	78.4	32 • 1	-71.5	45.9	-53.
95 .0	313.4	77.8	53.5	-56.5	63.8	-44.
96.0	312.6	88.6	60.0	-65.2	71.9	-51.7
97.0	325.0	107.9	88.4	-61.9	99•1	-42.8
98.0	324.9	108.9	89.2	-62.6	100.0	-43.3
99.0	325•6	109.6	90.4	-62.0	101.1	-42.5
190.0	326.3	114.5	95.2	-63.6	106.1	-43.1
101.0	334.8	53.9	48.8	-22.9	52.4	-12.6
102.0	27.3	17.8	15.8	8.1	13.8	11.1
103.0	122.3	51.4	-27.5	43.4	-35.7	37.0
104.0	138.5	69.6	-52 • 1	46 • 2	-60.4	34.7
105.0	145•2	94.5	-77•6	53.9	-86.9	37.1
106.0	146.1	103.6	-86.0	57.8	-95.9	39.2
107.0	143.5	108.9	-87.5	64.8	-98.8	45.8
108.0	148.4	108.9	-92•8	57.0	-102.4	37.1
109.9	156 • 8	103.0	-94.6	40.6	-100.8	20.7
110.0	160.6	91.1	-85.9	30.3	-90.2	12.3
111.0	156.7	83.0	-76.3	32.9	-81.4	16.8
112.0	145•1	70.7	-58.0	40.5	-65.0	28.0
113.0	131.7	66.4	-44.2	49.6	-53.3	39.7
114.0	113.0	69.2	-27.1	63.7	-39.4	56.9
115.0	105.2	72.2	-19.0	69.7	-32.7	64.4
116.0	191.0	71.9	-13.7	70.5	-27.7	66.3
117.0	95•7	70.9	-7.1	70.5	-21.2	67.6
118.0	89.5	73.7	0.6	73.7	-14.3	72.3
119.0	78.5	66.3	13.3	65.0	-0.1	66.3
120.0	66•9	65.8	25.9	60.5	13.1	64.5
121.0	61.6	59.8	28.5	52.6	17.3	57.3
122.0	41.2	53.9	40.5	35.5	32.5	42.9










03-27-00 AST

ALT	ITUDE	WIND HEADING	WIND VELOCITY		WIND COMPON		HETIC
	(KM)	(DEG)	(M/S)	N-S	E-W	N-S	E-w
	90.0	320.1	53.7	41.2	-34 • 4	47.3	-25.4
	91.0	324.6	52.7	43.0	-30.5	48.3	-21.2
	92.0	338.2	62.2	57.8	-23.1	61.3	-11.6
	93.0	337.4	71.6	66.1	-27.6	70.3	-13.7
	94.0	339.2	70.3	65.7	-25.0	69.4	-11.2
	95.0	349.5	58.1	57.2	-10.6	58.2	1.2
	96.0	1 • 1	43.9	43.9	0.9	42.8	9.7
	97.0	357.9	43.2	43.2	-1.6	42.6	7.2
	98.0	0.2	46.6	46.6	0 • 1	45.6	9.5
	99.0	355•8	49.8	49.7	-3.7	49 • 4	6.4
1	00.0	137.7	27.1	~20•0	18.2	-23.3	13.8
1	01.0	164.8	60.4	-58 • 3	15.8	-60.3	3.7
1	02.0	162.0	67.2	-64.0	20.6	-66.8	7.3
1	03.0	168.6	57.6	-66.3	13.3	-67.6	-0.4
1	04.0	176•2	74.7	-74.5	5•0	-74.0	-10.1
	05.0	184.3	78.8	-78.6	-6.0	-75.8	-21.7
	06.0	145.9	58.8	-48.7	32•9	-54.3	22.4
	07.0	129.0	66.2	-41.7	51.4	-51.2	41.9
	08.0	128.7	66.9	-41.9	52.2	-51.6	42.7
-	09.0	135.9	54.7	-39.3	38.1	-46.2	29.4
	10.0	139.5	29.2	-22.2	18.9	-25.6	14.0
	11.0	96 • 4	1.3	-0.1	1.2	-0.3	1.2
	12.0	352.7	12.2	12.1	-1.5	12.2	1.0
	13.0	352.5	18.5	18.4	-2 • 4	18.5	1.4
	14.0	333.8	38.4	34.5	-17.0	37.2	-9.7
	15.0	333•2	45 • 4	40.6	-20.5	43.9	-11.9
	16.0	331 • 1	58.2	51.0	-28 • 2	55.6	-17.3
	17.0	331.0	65.8	57.6	-31.9	62.9	-19.6
	18.0	335.7	71.1	64.8	-29.3	69.4	-15.6
	19.0	334 • 8	77.0	69.7	-32 • 8	74.9	-18.0
	20.0	337.1	80.8	74.4	-31.4	79.2	-15.7
	21.0	336 • 9	80.6 79.9	74•2 75•3	-31 • 7 -26 • 9	79•1 79•2	-16 · 1 -11 · 1
	22.0	340.3 339.8	86.1	80.8	-29•7	85.1	-12.8
į	634V	227 ♦ 0	00 • 7	0000	4701	0 ノ 🌣 メ	16.0









	WIND	WIND	WIND COMPONENTS (M/S))
ALTITUDE	HEADING	VELOCITY	GEOGRAPHIC			NETIC
(KM)	(DEG)	(M/S)	N-S	E-W	N-S	E-W
96.0	282 • 3	38.9	8 • 3	-38.0	15.8	
97.0	256 • 8	32.0	-7.3	-31.1	-0.9	-35.5 -31.9
98.0	226.2	38.7	-26.8	-27.9	-20.6	
99.0	205.1	119.1	-107.9	-50.5	-95.5	-32.7
100.0	206.7	131.2	-117.2	-58.9		-71.2
191.0	213.0	126.4	-106.0	-68.8	-102.9 -89.9	-81.4
102.0	219.5	111.4	-86.0	-70.9		-88.8
103.0	224.1	86.4	-62.1	-60.1	-69•9 -48•7	-80.8
104.0	225 • 8	58.3	-40.6	-41.8	-31·3	-71.4
105.0	224.9	34.7	-24.6	-24.5	-19•1	-49.1 -29.0
106.0	201.8	12.7	-11.8	-4.7	-10.6	
107.0	86•6	16.5	1.0	16.5	-10.6 -2.4	-7.0
108.0	78.4	36.2	7.3	35.5	0.0	16.4
109.0	72.3	41.7	12.7	39.7	4,4	36.2
110.0	64.7	44.2	18.9	39.9	10.5	41.4
111.0	52.1	44.3	27.2	35.0	19.6	42.9
112.0	35.5	52.5	42.7	30.5	35.7	39.8
113.0	23.2	64.3	59.1	25.4	52 • 8	38.5
114.0	22.9	79.9	73.6	31.1		36.8
115.0	33.9	89.3	74.1	49.8	65•8	45.3
116.9	18.9	67.3	63.7	21.8	62.5	63.7
117.0	4.9	49.0	48.8	4.2	58 • 0	34.2
				704	46.9	14.0


	WIND	WIND	WIND COMPONENTS (M/S))
4LTITUDE (KM)	HEADING	VELOCITY	GEOG	RAPHIC	MAGNETIC	
	(DEG)	(M/S)	N-S	E-W	N-S	E-W
101.0	220.2	121.7	-93.0	-78.5	-75.2	-95.7
102.0	222.8	99.2	-72.8	-67.4	-57.7	-80.7
103.0	226 • 1	70.5	-48.9	-50.8	-37.6	-59.6
104.0	224 • 3	56.8	-40.7	-39.7	-31.8	-47.1
105.0	224•6 223•8	43.4 19.5	-30.9 -14.0	-30.5 -13.5	-24 · 1 -11 · C	-36.1 -16.0
107.0	66 • 5 62 • 4	32.5 46.6	13.0 21.6	29 • 8 41 • 3	6 • 7 12 • 8	31.8
109.0 110.0 111.0	64•2 62•1 55•9	46.4 47.8 43.1	20.2 22.3 24.1	41.8 42.2 35.7	11.3 13.3 16.4	45.0 45.8 39.8
112.0	43 • 9 32 • 8	48.4 47.1	34.9 39.6	33.5 25.5	27 • 4 33 • 6	39.9 33.0
114.0	20.3 33.8	47.9 57.2	44.9 47.5	16.6 31.8	40.6 40.1	25.3 40.7
116.0 117.0	3419 26•6	59.9 46.4	49•1 41•5	34·3 20·7	41 • 2 36 • 5	43.5 28.7
118.0	347.1	24.6	24.0	-5.5	24.6	~0. 5


TRAIL A

ALTITUDE	WIND HEADING	WIND VELOCITY	GEOGR	VIND COMPONENTS (M/S)			
(11)				APHIC	MAGN	ETIC	
•	(DEG)	(M/S)	N-S	E-W	N-S	E-W	
116	10.9	54.4	53.4	10.2	50 •2	20.8	
111.0	19.9	65.4	61.5	22.3	5 5 • 7	34.3	
112.0	16.4	68.4	65.6	19.3	60.4	32.1	
113.0	359∙3	85 O	85.0	-1.0	83.5	16.2	
114.0	35%.2	85	85.3	-7.2	٠٠٥	10.2	












ALTITUDE	WIND HEADING	WIND VELOCITY	GEOG	WIND COMPO	NEHTS (M/S) NETIC
(KM)	(DEG)	(M/S)	N-S	E-W	N-S	
94.0	289.6	59.6	20.0	-56.2	07.9	E - #
95.0	312.0	61.4	41.1	~45•6		-51.0
96.0	323.1	65.8	52.6	-39.5	49.5 59.1	-35.4
97.0	337.6	74.7	69.1	-28.4	-	-28.1
98.€	348.7	85.1	83.5	-16.7	75.4	-13.9
99.0	356.9	99.8	99.6	-5.3	8:•/	2.5
100.0	3 • 8	119.4	119.1	8.0	98.6 115.0	14.5
191.0	5 • 8	133.3	132.6	13.6		31.5
102.0	11.3	141.9	139.2	27.7	127.1	40.1
163.0	15.5	135.2	130.3	36.1	130.7	51.2
104.0	20.5	133.6	125.1	47.0	120.3	61.7
105.0	34.9	163.2	84.7		113.3	71.3
106.0	44.5	114.7	81.9	59•1 80•3	71.0	75.0
167.0	63.0	132.7	60.3		64.0	95.2
168.0	7.0	176.8	28.5	118.2	35.2	127.9
109.0	92.0	100.7	-3.8	123.6	3.0	126.8
110.0	97.4	80.7	-10.4	106.5	-25.2	103.6
111.0	111.1	74.7	-27.0	80.0	-26.3	75.3
112.0	136.9	54.9	-40.1	69.7	-40.5	64.8
113.0	153.2	56.1		37.6	-46.9	28.7
114.0	159.4	47.2	-50.0 -44.2	25.3	-54.1	14.7
115.0	166.5		-44.2	16.6	-46.6	7.3
116.0	173.4	41.9 37.3	-40.7	9 • 8	-41.8	1.4
117.0	183.8	34.0	-37.1	4 • 3	-37.2	-3.3
118.0	193.5	34.0 30.7	~33.9	-2 • 2	-32.8	-9.0
119.0	204.5	26.7	-29.9	-7.2	-27.8	-13.1
126.0	212.0	27.3	-24.3	-11.1	-21.6	-15.8
121.0	220.6		-23.1	-14.5	-19.7	-18.9
122.0	221.7	26.0 27.1	-19.7	-16.9	-15.9	-20.5
123.0	216.5		-20.3	-18.0	-16.2	-2i.,
124.0	214.1	26.6	-21.4	-15.8	-17.8	-17.8
125.0	207.5	26.9	-22.3	-15.1	-18.8	-19.3
126.0	190.9	² 4.0 19.9	-21.3	-11.1	-18.6	-15.2
127.0	170.2		-19.6	-3 • 8	-18.4	-7.7
128.0	154.5	20•4 22•0	-20.1	3 • 5	-20.4	-0.6
129.0	135.5		-19.9	9.5	-21.4	5.3
130.0	123.7	25.5	-18.2	17.8	-21.4	13.8
	16301	31.5	-17.5	26.2	-22.4	22.1







Security Classification DCCUMENT CONTI	ROL DATA - R & D
	nustation must be entered when the overall report is classified)
Space Instruments Research, Inc.	Unclassified 26. GROUP
TPPER ATTOSPHERE MINDS FROT GUT-LAU 17-25 FEDRUARY 1966)	CHED VERTICAL PRODES (BIRBADOS,
4. DESCRIPTIVE NOTES (Type of report end Inclus ve dites)	
5. AUTHO (5) (i not no occupanted)	
Robert L. Fuller	•
6. REPORT DATE	7a. TOTAL NO. OF PAGES 7b. NO OF REFS
January 1967	54 37
ES. CONTRACT OR GRANT NOUN - () - (1 - (1 - (1 - (1 - (1 - (1 - (1	DJ. ORIGINATON'S REPORT NUMBER(S)
ь. Расочест но. RDTE 1V014501053C	EFL Renort 169 Penort 5
c.	9b. OTHER REPORT NO(5) (Any other n. Store that may be easilized this report)
d.	
16. DISTRIBUTION STATEMENT	
This document has been approved for pub is unlimited.	lic release and sale; its distribution
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY
	Commanding Officer U.S. Army Fallistic Research Laboratories
	Aberdeen Proving Ground, Md. 21005
13. ADSTRACT	2000
on the night of 23-24 February produced between 87km and 131km by	1966, six luminous trails were
from projectiles fired from a sport	bhore sixteen-inch aun located on
the West Indian island of Barbados	(57.5°U, 13.1°U). Single trails
were also produced on the nights of	1/ February 1966 and 25 February
1966. These trails were photograph	ed from neighboring islands and
analyzed to yield wind profiles. T wind data for all eight trails toge	ther with plots versus altitude of
wind data for all eight trails toge wind components, wind speed, and wi	nd heading.
with the composition of the contract of the co	
	•
•	
· ·	
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Augus (1975) deuts au Statt (S. Statte) au (1975) attaut at the Courage (1974) at the Courage (1975) attaut attaut at the Courage (1975) attaut attaut at the Courage (1975) attaut at

UnCLASSIFIED
Security Classification

UNCLASSIFIED
Security Classification

n kenegarang dinang melilipin dinang belang dinang kenegarang belang dinang belang dinang belang belang belang Ak	KEY NONDS		L'NK A LIN			LIR	K C
ma hadia variandridhi yawana ayddi diri		ROLE	WT	ROLE	V: T	ROLE	WT
TRP igh Altitude onospheric Vi	Pescarch Project nds	,					
			•				
	·						
•							
	·						
		•					
	•						

	ASS		
	with C		*10"